

Representations of quantum affine Lie algebras

Vyacheslav Futorny

Shenzhen International Center for Mathematics
Southern University of Science and Technology

Joint results with Xingpeng Liu, Juan Camilo Arias and Kailash Misra

Linköping - 2026

Affine Lie algebras

\mathfrak{g} a simple finite-dimensional Lie algebra

Loop algebra $\tilde{\mathfrak{g}} = \mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]$, with the Lie bracket
 $[x \otimes t^n, y \otimes t^m] = [x, y] \otimes t^{m+n}$

Non-twisted affine Kac-Moody algebra is the universal central extension $\hat{\mathfrak{g}} = \tilde{\mathfrak{g}} \oplus \mathbb{C}c \oplus \mathbb{C}d$,

$$[x \otimes t^n, y \otimes t^m] = [x, y] \otimes t^{m+n} + n(x, y)\delta_{n+m,0}c,$$

$d : \hat{\mathfrak{g}} \rightarrow \hat{\mathfrak{g}}$ degree derivation, $d(x \otimes t^n) = n(x \otimes t^n)$, $d(c) = 0$

$\hat{\mathfrak{h}} = \mathfrak{h} \otimes_{\mathbb{C}} \mathbb{C}1 \oplus \mathbb{C}c \oplus \mathbb{C}d$ **Cartan subalgebra**

Borel and parabolic subalgebras

◇ A subset $P \subset \Delta$ is a **quase partition** if:

a) $P \cap (-P) = \emptyset$ and $P \cup (-P) = \Delta$

b) Let \mathcal{B}_P a Lie subalgebra of $\widehat{\mathfrak{g}}$ generated by $\widehat{\mathfrak{h}}$ and the root spaces $\widehat{\mathfrak{g}}_\alpha$ with $\alpha \in P$. Then for any root α of \mathcal{B}_P we have $\alpha \in P$

◇ \mathcal{B}_P describes (almost all) Borel subalgebras of $\widehat{\mathfrak{g}}$. There exists a finite number conjugacy classes, (roughly) parameterized by the parabolic subalgebras of \mathfrak{g} (**Jacobsen-Kac; VF**)

Example

- **standard** Borel subalgebra with the partition $P = \Delta^+$
- **natural** Borel subalgebra \mathcal{B}_{nat} with the partition

$$P_{\text{nat}} = \{\alpha + k\delta \mid \alpha \in \Delta^+(\mathfrak{g}), k \in \mathbb{Z}\} \cup \{n\delta \mid n \geq 0\}$$

- each function $\phi : \mathbb{N} \rightarrow \mathbb{Z}_2$ defines a Borel subalgebra $\mathcal{B}_{\text{nat}}^\phi$ with quase partition $\{\alpha + k\delta \mid \alpha \in \Delta^+, k \in \mathbb{Z}\} \cup \{n\delta \mid n \in \mathbb{N}, \phi(n) = 1\} \cup \{-m\delta \mid m \in \mathbb{N}, \phi(m) = 0\}$

Closed partitions in $\hat{\mathfrak{sl}}_2$

Standard partition:

$$\begin{array}{cccccccc} \cdots & -\alpha_0 - \delta & -\alpha_0 & \alpha_1 & \alpha_1 + \delta & \alpha_1 + 2\delta & \alpha_1 + 3\delta & \cdots \\ \cdots & -2\delta & -\delta & & \delta & 2\delta & 3\delta & \cdots \\ \cdots & -\alpha_1 - 2\delta & -\alpha_1 - \delta & -\alpha_1 & \alpha_0 & \alpha_0 + \delta & \alpha_0 + 2\delta & \cdots \end{array}$$

Closed partitions in $\hat{\mathfrak{sl}}_2$

Standard partition:

$$\begin{array}{cccccccc} \cdots & -\alpha_0 - \delta & -\alpha_0 & \alpha_1 & \alpha_1 + \delta & \alpha_1 + 2\delta & \alpha_1 + 3\delta & \cdots \\ \cdots & -2\delta & -\delta & & \delta & 2\delta & 3\delta & \cdots \\ \cdots & -\alpha_1 - 2\delta & -\alpha_1 - \delta & -\alpha_1 & \alpha_0 & \alpha_0 + \delta & \alpha_0 + 2\delta & \cdots \end{array}$$

Natural partition:

$$\begin{array}{cccccccc} \cdots & -\alpha_0 - \delta & -\alpha_0 & \alpha_1 & \alpha_1 + \delta & \alpha_1 + 2\delta & \alpha_1 + 3\delta & \cdots \\ \cdots & -2\delta & -\delta & & \delta & 2\delta & 3\delta & \cdots \\ \cdots & -\alpha_1 - 2\delta & -\alpha_1 - \delta & -\alpha_1 & \alpha_0 & \alpha_0 + \delta & \alpha_0 + 2\delta & \cdots \end{array}$$

◇ A **closed** subset $P \subset \Delta$ such that $P \cup (-P) = \Delta$ defines a **parabolic subalgebra** $\mathcal{P} \subset \widehat{\mathfrak{g}}$

◇ A parabolic subalgebra $\mathcal{P} \subset \widehat{\mathfrak{g}}$ contains a Borel subalgebra:

a) **type I** (contains the standard Borel):

$\mathcal{P} = \mathfrak{l} \oplus \mathfrak{u}$, where \mathfrak{l} is finite-dimensional reductive Lie algebra;

b) **type II** (contains one of Borel subalgebras $\mathcal{B}_{\text{nat}}^\phi$):

$\mathcal{P} = \mathfrak{l} \oplus \mathfrak{u}$, where \mathfrak{l} is infinite-dimensional Lie subalgebra

Parabolic induction

◇ Borel subalgebra $\mathcal{B} \rightsquigarrow$ Verma type module

$$M_{\mathcal{B}}(\lambda) = U(\widehat{\mathfrak{g}}) \otimes_{U(\mathcal{B})} \mathbb{C}$$

◇ Parabolic subalgebra $\mathcal{P} \rightsquigarrow$ generalized Verma type module

$$M_{\mathcal{P}}(N) = U(\widehat{\mathfrak{g}}) \otimes_{U(\mathcal{P})} N$$

Theorem (F., Kashuba et al, 2016-2023)

Let $\mathcal{P} \subset \widehat{\mathfrak{g}}$ be a parabolic subalgebra of type II, N is a nice simple \mathcal{P} -module with $c \neq 0$ Then $M_{\mathcal{P}}(N)$ is a simple $\widehat{\mathfrak{g}}$ -module

Imaginary Verma modules

For the natural closed partition

$$S = \{\alpha + n\delta \mid \alpha \in \Delta^+(\mathfrak{g}), n \in \mathbb{Z}\} \cup \{k\delta \mid k > 0\}$$

consider the following subalgebras of $\hat{\mathfrak{g}}$: $\mathfrak{g}_{\pm}(S) = \sum_{\alpha \in S} \hat{\mathfrak{g}}_{\pm\alpha}$.

For example, for $\mathfrak{g} = \mathfrak{sl}(2)$ we have

► $\mathfrak{g}_+(S)$ is the subalgebra generated by $e(k) = \begin{pmatrix} 0 & t^k \\ 0 & 0 \end{pmatrix}$, ($k \in \mathbb{Z}$) and

$$h(l) = \begin{pmatrix} t^l & 0 \\ 0 & -t^l \end{pmatrix}, \quad (l \in \mathbb{Z}_{>0}).$$

► $\mathfrak{g}_-(S)$ is the subalgebra generated by $f(k) = \begin{pmatrix} 0 & 0 \\ t^k & 0 \end{pmatrix}$, ($k \in \mathbb{Z}$) and

$$h(-l) = \begin{pmatrix} t^{-l} & 0 \\ 0 & -t^{-l} \end{pmatrix}, \quad (l \in \mathbb{Z}_{>0}),$$

$$\hat{\mathfrak{g}} = \mathfrak{g}_-(S) \oplus \hat{\mathfrak{h}} \oplus \mathfrak{g}_+(S),$$

and

$$U(\hat{\mathfrak{g}}) \cong U(\mathfrak{g}_-(S)) \otimes U(\hat{\mathfrak{h}}) \otimes U(\mathfrak{g}_+(S))$$

A $U(\hat{\mathfrak{g}})$ -module V is called a *weight* module if $V = \bigoplus_{\mu \in \hat{\mathfrak{h}}^*} V_{\mu}$, where

$$V_{\mu} = \{v \in V \mid h \cdot v = \mu(h)v, c \cdot v = \mu(c)v, d \cdot v = \mu(d)v\}.$$

A $U(\hat{\mathfrak{g}})$ -module V is called an *S-highest weight module* with highest weight λ if there is a non-zero $v_{\lambda} \in V$ such that


- ▶ $u^+ \cdot v_{\lambda} = 0$ for all $u^+ \in \mathfrak{g}_+(S) \setminus \mathbb{C}^*$,
- ▶ $h \cdot v_{\lambda} = \lambda(h)v_{\lambda}$, $c \cdot v_{\lambda} = \lambda(c)v_{\lambda}$, $d \cdot v_{\lambda} = \lambda(d)v_{\lambda}$,
- ▶ $V = U(\hat{\mathfrak{g}}) \cdot v_{\lambda} = U(\mathfrak{g}_-(S)) \cdot v_{\lambda}$.


For $\lambda \in \hat{\mathfrak{h}}^*$, let $I_S(\lambda)$ denote the ideal of $U(\widehat{\mathfrak{sl}(2)})$ generated by $e(k)$ ($k \in \mathbb{Z}$), $h(l)$ ($l > 0$), $h - \lambda(h)1$, $c - \lambda(c)1$, $d - \lambda(d)1$. Then the *imaginary Verma module* of $\widehat{\mathfrak{sl}(2)}$ with highest weight λ is defined as

$$M_S(\lambda) = U(\widehat{\mathfrak{sl}(2)})/I_S(\lambda)$$


Theorem

- (i) $M_S(\lambda)$ is a $U(\mathfrak{g}_-(S))$ -free module of rank 1 generated by the S -highest weight vector $1 \otimes 1$ of weight λ .
- (ii) $\dim M_S(\lambda)_\lambda = 1$; $0 < \dim M_S(\lambda)_{\lambda - k\delta} < \infty$ for any integer $k > 0$; if $\mu \neq \lambda - k\delta$ for any integer $k \geq 0$ and $M_S(\lambda)_\mu \neq 0$, then $\dim M(\lambda)_\mu = \infty$.
- (iii) $M_S(\lambda)$ has a unique maximal submodule.
- (iv) $M_S(\lambda)$ is irreducible if and only if $\lambda(c) \neq 0$.
- (v) Let $\lambda(c) = 0$ then $(U(\sum_{k \in \mathbb{Z} \setminus \{0\}} \hat{\mathfrak{g}}_{k\delta} \oplus \mathbb{C}c) \setminus \{\mathbb{C}\}) \cdot 1$ is a proper submodule of $M_S(\lambda)$. Moreover, it is maximal if and only if $\lambda(h) \neq 0$





Let $\text{Rep}U_q(\mathfrak{g})$ be the category of finite-dimensional $U_q(\mathfrak{g})$ -modules



- Recall that a simple weight Harish-Chandra \mathfrak{g} -module V ($\dim V_\lambda < \infty$ for any $\lambda \in \mathfrak{h}^*$) which cannot be induced from a module over any proper parabolic subalgebra of \mathfrak{g} is called *cuspidal*. It is known that such modules exist only when \mathfrak{g} consists of simple components of type A or type C .

Theorem (F-Tsylke, 2001)

Let V be a simple Harish-Chandra $\hat{\mathfrak{g}}$ -module with a nonzero action of c . Then $V \simeq L_{k,\mathfrak{p}}(N)$, for some parabolic subalgebra $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u}$ of \mathfrak{g} and a simple cuspidal \mathfrak{l} -module $N \simeq \otimes D_{S_i}^{\mu_i} L(\lambda_i)$ for some sets of roots S_i 's, $\mu_i \in \mathfrak{h}^$ and simple λ_i -highest weight \mathfrak{l}_i -modules $L(\lambda_i)$, where \mathfrak{l}_i 's are simple components of \mathfrak{l} of type A and C .*

- Let $\check{U}_q := U_q \otimes_{\mathbb{C}} \mathbb{k}$, where \mathbb{k} is the algebraic closure of $\mathbb{C}(q)$.

Theorem (V.F., X.Liu, 2025)

The algebra \check{U}_q admits cuspidal modules if and only if the underlying semisimple Lie algebra \mathfrak{g} consists of simple components of type A, B or C (in contrast, for $U_{\mathbb{Q}(q)}$ cuspidal modules of type B do not exist).

Remark: The result holds not only for an indeterminate q but also for any specialization $q \rightarrow \xi$, where $\xi \in \mathbb{C}$ is transcendental.

Example (X.Liu)

Consider the quantum group $U_q(\mathfrak{g})$ of type $B_2(= C_2)$ with the Cartan matrix

$$(a_{ij}) = \begin{pmatrix} 2 & -1 \\ -2 & 2 \end{pmatrix},$$

and generators $E_i, F_i, K_i^{\pm 1}, i = 1, 2$. Set $q_1 = q^2, q_2 = q$.

Define V as the vector space spanned by $|m_1, m_2\rangle$ for $m_1, m_2 \in \mathbb{Z}_{\geq 0}$. There exists a $U_q(\mathfrak{g})$ -module structure on V :

$$\begin{aligned} K_1 \cdot |m_1, m_2\rangle &= q_1^{-m_1+m_2} |m_1, m_2\rangle, & K_2 \cdot |m_1, m_2\rangle &= \sqrt{-1} q_2^{-2m_2-1} |m_1, m_2\rangle, \\ E_1 \cdot |m_1, m_2\rangle &= [m_1]_{q_1} |m_1 - 1, m_2 + 1\rangle, & E_2 \cdot |m_1, m_2\rangle &= \sqrt{-1} \frac{q_2 + q_2}{q_2 - q_2} [m_2]_{q_1} |m_1, m_2 - 1\rangle, \\ F_1 \cdot |m_1, m_2\rangle &= [m_2]_{q_1} |m_1 + 1, m_2 - 1\rangle, & F_2 \cdot |m_1, m_2\rangle &= |m_1, m_2 + 1\rangle. \end{aligned}$$

Example (X.Liu)

Consider the quantum group $U_q(\mathfrak{g})$ of type $B_2(= C_2)$ with the Cartan matrix

$$(a_{ij}) = \begin{pmatrix} 2 & -1 \\ -2 & 2 \end{pmatrix},$$

and generators $E_i, F_i, K_i^{\pm 1}, i = 1, 2$. Set $q_1 = q^2, q_2 = q$.

Define V as the vector space spanned by $|\mathbf{m}_1, \mathbf{m}_2\rangle$ for $m_1, m_2 \in \mathbb{Z}_{\geq 0}$. There exists a $U_q(\mathfrak{g})$ -module structure on V :

$$\begin{aligned} K_1 \cdot |\mathbf{m}_1, \mathbf{m}_2\rangle &= q_1^{-m_1+m_2} |\mathbf{m}_1, \mathbf{m}_2\rangle, & K_2 \cdot |\mathbf{m}_1, \mathbf{m}_2\rangle &= \sqrt{-1} q_2^{-2m_2-1} |\mathbf{m}_1, \mathbf{m}_2\rangle, \\ E_1 \cdot |\mathbf{m}_1, \mathbf{m}_2\rangle &= [m_1]_{q_1} |\mathbf{m}_1 - 1, \mathbf{m}_2 + 1\rangle, & E_2 \cdot |\mathbf{m}_1, \mathbf{m}_2\rangle &= \sqrt{-1} \frac{q_2 + q_2}{q_2 - q_2} [m_2]_{q_1} |\mathbf{m}_1, \mathbf{m}_2 - 1\rangle, \\ F_1 \cdot |\mathbf{m}_1, \mathbf{m}_2\rangle &= [m_2]_{q_1} |\mathbf{m}_1 + 1, \mathbf{m}_2 - 1\rangle, & F_2 \cdot |\mathbf{m}_1, \mathbf{m}_2\rangle &= |\mathbf{m}_1, \mathbf{m}_2 + 1\rangle. \end{aligned}$$

This module has a nice property that it is infinite dimensional, the “minimal” bounded weight module for $U_q(\mathfrak{g})$.

The quantum group $U_q(\hat{\mathfrak{g}})$

We denote by $U_q(\hat{\mathfrak{g}})$ the **quantum affine algebra** associated to $\hat{\mathfrak{g}}$: the associative unital $\mathbb{C}(q^{1/2})$ -algebra with generators $E_i, F_i, K_\alpha, \gamma^{\pm 1/2}, D^{\pm 1}$ for $0 \leq i \leq N$, $\alpha \in Q$ and defining relations:

$$DD^{-1} = D^{-1}D = K_\alpha K_{-\alpha} = K_{-\alpha} K_\alpha = \gamma^{1/2} \gamma^{-1/2} = \gamma^{-1/2} \gamma^{1/2} = 1$$

$$[\gamma^{\pm 1/2}, U_q(\hat{\mathfrak{g}})] = [D, K_{\pm\alpha}] = [K_\alpha, K_\beta] = 0$$

$$(\gamma^{\pm 1/2})^2 = K_{\pm\delta}$$

$$E_i F_j - F_j E_i = \delta_{ij} \frac{K_i - K_i^{-1}}{q_i - q_i^{-1}}$$

$$K_\alpha E_i K_{-\alpha} = q^{(\alpha|\alpha_i)} E_i, \quad K_\alpha F_i K_{-\alpha} = q^{-(\alpha|\alpha_i)} F_i$$

$$DE_i D^{-1} = q^{\delta_{i,0}} E_i, \quad DF_i D^{-1} = q^{-\delta_{i,0}} F_i$$

$$\sum_{s=0}^{1-a_{ij}} (-1)^s E_i^{(1-a_{ij}-s)} E_j E_i^{(s)} = \sum_{s=0}^{1-a_{ij}} (-1)^s F_i^{(1-a_{ij}-s)} F_j F_i^{(s)} = 0, \quad i \neq j$$

where $q_i = q^{d_i}$, $[n]_i = \frac{q_i^n - q_i^{-n}}{q_i - q_i^{-1}}$, $[n]_i! = [n]_i [n-1]_i \cdots [2]_i [1]_i$, $K_i = K_{\alpha_i}$,

$E_i^{(s)} = E_i^s / [s]_i!$ and $F_i^{(s)} = F_i^s / [s]_i!$

Similar to the loop space realization of $\hat{\mathfrak{g}}$ we have the Drinfel'd realization: the generators are $x_{ir}^{\pm 1}, h_{is}, K_i^{\pm 1}, \gamma^{\pm 1/2}, D^{\pm 1}$ for $i \in I_0 = \{1, \dots, n\}$, $r, s \in \mathbb{Z}$ and $s \neq 0$ subject to the relations:

$$\begin{aligned}
 D^{\pm 1} D^{\mp 1} &= K_i^{\pm 1} K_i^{\mp 1} = \gamma^{\pm 1/2} \gamma^{\mp 1/2} = 1 \\
 [\gamma^{\pm 1/2}, U_q(\hat{\mathfrak{g}})] &= [D, K_i^{\pm 1}] = [K_i, K_j] = [K_i, h_{js}] = 0 \\
 Dh_{ir} D^{-1} &= q^r h_{ir}, \quad Dx_{ir}^{\pm} D^{-1} = q^r x_{ir}^{\pm} \\
 K_i x_{jr}^{\pm} K_i^{-1} &= q^{\pm(\alpha_i | \alpha_j)} x_{jr}^{\pm} \\
 [h_{ik}, h_{jl}] &= \delta_{k,-l} \frac{1}{k} [ka_{ij}]_i \frac{\gamma^k - \gamma^{-k}}{q_j - q_j^{-1}} \\
 [h_{ik}, x_{jl}^{\pm}] &= \pm \frac{1}{k} [ka_{ij}]_i \gamma^{\mp |k|/2} x_{j,k+l}^{\pm} \\
 x_{i,k+1}^{\pm} x_{jl}^{\pm} - q^{\pm(\alpha_i | \alpha_j)} x_{jl}^{\pm} x_{i,k+1}^{\pm} &= q^{\pm(\alpha_i | \alpha_j)} x_{ik}^{\pm} x_{j,l+1}^{\pm} - x_{j,l+1}^{\pm} x_{ik}^{\pm} \\
 [x_{ik}^+, x_{jl}^-] &= \delta_{ij} \frac{1}{q_i - q_i^{-1}} (\gamma^{(k-l)/2} \psi_{i,k+l} - \gamma^{(l-k)/2} \phi_{i,k+l})
 \end{aligned}$$

where

$$\begin{aligned}
 \sum_{k=0}^{\infty} \psi_{ik} z^k &= K_i \exp \left((q_i - q_i^{-1}) \sum_{l>0} h_{il} z^l \right) \\
 \sum_{k=0}^{\infty} \phi_{i,-k} z^{-k} &= K_i^{-1} \exp \left(-(q_i - q_i^{-1}) \sum_{l>0} h_{i,-l} z^{-l} \right)
 \end{aligned}$$

and for $i \neq j$,

$$\text{Sym}_{k_1, \dots, k_{1-a_{ij}}} \sum_{r=0}^{1-a_{ij}} (-1)^r \begin{bmatrix} 1 - a_{ij} \\ r \end{bmatrix}_i x_{ik_1}^\pm \cdots x_{ik_r}^\pm x_{ij}^\pm x_{ik_{r+1}}^\pm \cdots x_{ik_{1-a_{ij}}}^\pm = 0.$$

If we consider the following generating functions

$$\phi_i(u) = \sum_{p \in \mathbb{Z}} \phi_{ip} u^{-p}, \quad \psi_i(u) = \sum_{p \in \mathbb{Z}} \psi_{ip} u^{-p}, \quad x_i^\pm(u) = \sum_{p \in \mathbb{Z}} x_{ip}^\pm u^{-p}$$

the defining relations become:

$$\begin{aligned} [\phi_i(u), \phi_j(v)] &= [\psi_i(u), \psi_j(v)] = 0 \\ \phi_i(u)\psi_j(v)\phi_i(u)^{-1}\psi_j(v)^{-1} &= g_{ij}(uv^{-1}\gamma^1)/g_{ji}(uv^{-1}\gamma) \\ \phi_i(u)x_j^\pm(v)\phi_i(u)^{-1} &= g_{ij}(uv^{-1}\gamma^{\mp 1/2})^{\pm 1}x_j^\pm(v) \\ \psi_i(u)x_j^\pm(v)\psi_i(u)^{-1} &= g_{ji}(vu^{-1}\gamma^{\mp 1/2})^{\mp 1}x_j^\pm(v) \\ (u - q^{\pm(\alpha_i|\alpha_j)v})x_i^\pm(u)x_j^\pm(v) &= (q^{\pm(\alpha_i|\alpha_j)u-v})x_j^\pm(v)x_i^\pm(u) \\ [x_i^+(u), x_j^-(v)] &= \delta_{ij}(q_i - q_i^{-1})(\delta(u/v\gamma)\psi_i(v\gamma^{1/2}) - \delta(u\gamma/v)\phi_i(u\gamma^{1/2})) \end{aligned}$$

where $g_{ij}(t) = g_{ij,q}(t)$ is the Taylor expansion at $t = 0$ of the function $(q^{(\alpha_i|\alpha_j)t} - 1)/(t - q^{(\alpha_i|\alpha_j)})$ and $\delta(z) = \sum_{k \in \mathbb{Z}} z^k$.

Conjecture [V.F., X.Liu, 2026]:

Let V_q be a simple Harish-Chandra $U_q(\hat{\mathfrak{g}})$ -module with a nonzero action of c . Then $V_q \simeq L_{k,p}(N_q)$, for some parabolic subalgebra $\mathfrak{p} = \mathfrak{l} \oplus \mathfrak{u}$ of \mathfrak{g} and a simple cuspidal $U_q(\mathfrak{l})$ -module $N \simeq \otimes D_{S_i}^{\mu_i} L_q(\lambda_i)$ for some sets of roots S_i 's, $\mu_i \in \mathfrak{h}^*$ and simple λ_i -highest weight \mathfrak{l}_i -modules $L_q(\lambda_i)$, where \mathfrak{l}_i 's are simple components of \mathfrak{l} of type A , B and C .

Quantum imaginary Verma modules

For the natural closed partition

$$S = \{\alpha + n\delta \mid \alpha \in \Delta^+(\mathfrak{g}), n \in \mathbb{Z}\} \cup \{k\delta \mid k > 0\}$$

consider the following subalgebras of $U_q(\hat{\mathfrak{g}})$:

- ▶ $U_q^+(S)$ generated by x_{ik}^+, h_{il} for $i \in I_0, k \in \mathbb{Z}$ and $l > 0$.
- ▶ $U_q^-(S)$ generated by $x_{ik}^-, h_{i,-l}$ for $i \in I_0, k \in \mathbb{Z}$ and $l > 0$.
- ▶ $U_q^0(S)$ generated by $K_i^{\pm 1}, \gamma^{\pm 1/2}, D^{\pm 1}$ for $i \in I_0$.

Let P be the integral weight lattice of $\hat{\mathfrak{g}}$, $\lambda \in P$. A weight module V of $U_q(\hat{\mathfrak{g}})$ is called an **S-highest weight module** with highest weight λ if there is a non zero vector $v \in V$ of weight λ such that $u^+v = 0$ for all $u^+ \in U_q^+(S) \setminus \mathbb{C}(q^{1/2})$ and $V = U_q(\hat{\mathfrak{g}})v$

Consider the Borel subalgebra B_q of $U_q(\hat{\mathfrak{g}})$ generated by $U_q^+(S) \cup U_q^0(\hat{\mathfrak{g}})$, and a one dimensional B_q -module $\mathbb{C}(q^{1/2})_\lambda$ with a generator $\mathbf{1}$, on which $U_q^+(S)$ acts trivially and $K_i^{\pm 1} \mathbf{1} = q^{\pm \lambda(h_i)} \mathbf{1}$, $i \in I_0$, $\gamma^{\pm 1/2} \mathbf{1} = q^{\pm \lambda(c)/2} \mathbf{1}$ and $D^{\pm 1} \mathbf{1} = q^{\pm \lambda(d)} \mathbf{1}$.

The **imaginary Verma module** $M_q(\lambda)$ of weight $\lambda \in P$ is defined as

$$M_q(\lambda) = M_{q,S}(\lambda) := U_q(\hat{\mathfrak{g}}) \otimes_{B_q} \mathbb{C}(q^{1/2})_\lambda$$

Consider the Borel subalgebra B_q of $U_q(\hat{\mathfrak{g}})$ generated by $U_q^+(S) \cup U_q^0(\hat{\mathfrak{g}})$, and a one dimensional B_q -module $\mathbb{C}(q^{1/2})_\lambda$ with a generator $\mathbf{1}$, on which $U_q^+(S)$ acts trivially and $K_i^{\pm 1} \mathbf{1} = q^{\pm \lambda(h_i)} \mathbf{1}$, $i \in I_0$, $\gamma^{\pm 1/2} \mathbf{1} = q^{\pm \lambda(c)/2} \mathbf{1}$ and $D^{\pm 1} \mathbf{1} = q^{\pm \lambda(d)} \mathbf{1}$.

The **imaginary Verma module** $M_q(\lambda)$ of weight $\lambda \in P$ is defined as

$$M_q(\lambda) = M_{q,S}(\lambda) := U_q(\hat{\mathfrak{g}}) \otimes_{B_q} \mathbb{C}(q^{1/2})_\lambda$$

Theorem [F., Grishkov, Melville, 2005]

$M_q(\lambda)$ is irreducible if and only if $\lambda(c) \neq 0$.

- Let $\lambda(c) = 0$. Denote by $J^q(\lambda)$ the left ideal of $U_q(\hat{\mathfrak{g}})$ generated by x_{ik}^+, h_{il} for $i \in I_0, k, l \in \mathbb{Z}, l \neq 0$ and $K_i^{\pm 1} - q^{\pm \lambda(h_i)}, \gamma^{\pm 1/2} - 1$ and $D^{\pm 1} - q^{\pm \lambda(d)}$. Set

$$\tilde{M}_q(\lambda) = U_q(\hat{\mathfrak{g}})/J^q(\lambda)$$

It is a quotient of $M_q(\lambda)$, we call it **reduced imaginary Verma module**.

- Let $\lambda(c) = 0$. Denote by $J^q(\lambda)$ the left ideal of $U_q(\hat{\mathfrak{g}})$ generated by x_{ik}^+ , h_{il} for $i \in I_0$, $k, l \in \mathbb{Z}$, $l \neq 0$ and $K_i^{\pm 1} - q^{\pm \lambda(h_i)}$, $\gamma^{\pm 1/2} - 1$ and $D^{\pm 1} - q^{\pm \lambda(d)}$. Set

$$\tilde{M}_q(\lambda) = U_q(\hat{\mathfrak{g}})/J^q(\lambda)$$

It is a quotient of $M_q(\lambda)$, we call it **reduced imaginary Verma module**.

Theorem [F., Grishkov, Melville]

Let $\lambda \in P$ such that $\lambda(c) = 0$. Then

- ▶ $\tilde{M}_q(\lambda)$ is irreducible if and only if $\lambda(h_i) \neq 0$ for all $i \in I_0$.
- ▶ $\tilde{M}_q(\lambda) = \bigoplus_{\substack{i_1, \dots, i_r \\ k_1, \dots, k_r}} \mathbb{C}(q^{1/2}) x_{i_1 k_1}^- \cdots x_{i_r k_r}^- v_\lambda$, where v_λ stands for the generator of the module.

- $U_q \cong U_q^-(S) \otimes U_q^0(S) \otimes U_q^+(S)$ and a PBW basis

Kashiwara algebra

- Consider the subalgebra \mathcal{N}_q^- of $U_q(\hat{\mathfrak{g}})$ generated by $\gamma^{\pm 1/2}$ and x_{ij}^- for $l \in \mathbb{Z}$, $i \in I_0$.

Set $\bar{P}^{j_1, \dots, j_k} = x_{j_1}^-(v_1) \cdots x_{j_k}^-(v_k)$,

$\bar{P}_l^{j_1, \dots, j_k} = x_{j_1}^-(v_1) \cdots x_{j_{l-1}}^-(v_{j_{l-1}}) x_{j_{l+1}}^-(v_{j_{l+1}}) \cdots x_{j_k}^-(v_k)$

and denote

$$G_{il} := \delta_{i, j_l} \prod_{m=1}^{l-1} g_{i, j_m, q^{-1}}(v_{j_m}/v_l), \quad G_{i1} = \delta_{i, j_1}$$

- We define operators $\Omega_i(k) : \mathcal{N}_q^- \longrightarrow \mathcal{N}_q^-$ for $k \in \mathbb{Z}$ in terms of the generating functions $\Omega_i(u) = \sum_{l \in \mathbb{Z}} \Omega_i(l) u^{-l}$:

$$\Omega_i(u)(\bar{P}^{j_1, \dots, j_k}) = \sum_{l=1}^k G_{il} \bar{P}_l^{j_1, \dots, j_k} \delta(u/v_l \gamma),$$

$\Omega_i(u)(1) = 0$. Also consider left multiplication operators $x_{im}^- : \mathcal{N}_q^- \longrightarrow \mathcal{N}_q^-$. The Ω -operators and the x^- -operators satisfy the identities:

$$q^{(\alpha_i|\alpha_j)} \gamma \Omega_j(m) x_{i,n+1}^- - \Omega_j(m+1) x_{in}^- = (q^{(\alpha_i|\alpha_j)} \gamma - 1) \delta_{ij} \delta_{m,-n-1} + \gamma x_{i,n+1}^- \Omega_j(m) - q^{(\alpha_1|\alpha_j)} x_{in}^- \Omega_j(m+1).$$

and

$$\Omega_j(k) x_{im}^- = \delta_{ij} \delta_{k,-m} \gamma^k + \sum_{r \geq 0} g_{i,j,q^{-1}}(r) x_{i,m+r}^- \Omega_j(k-r) \gamma^r.$$

- Define the **Kashiwara algebra** \mathcal{K}_q : the $\mathbb{C}(q^{1/2})$ -algebra with generators $\Omega_j(m), x_i^-(n), \gamma^{\pm 1/2}$ for $m, n \in \mathbb{Z}, 1 \leq i, j \leq N$, where $\gamma^{\pm 1/2}$ are central, $\gamma^{\pm 1/2} \gamma^{\mp 1/2} = 1$ and

$$q^{(\alpha_i|\alpha_j)} \gamma \Omega_j(m) x_{i,n+1}^- - \Omega_j(m+1) x_{in}^- =$$

$$(q^{(\alpha_i|\alpha_j)} \gamma - 1) \delta_{ij} \delta_{m,-n-1} + \gamma x_{i,n+1}^- \Omega_j(m) - q^{(\alpha_i|\alpha_j)} x_{in}^- \Omega_j(m+1)$$

$$q^{(\alpha_i|\alpha_j)} \Omega_i(k+1) \Omega_j(l) - \Omega_j(l) \Omega_i(k+1) =$$

$$\Omega_i(k) \Omega_j(l+1) - q^{(\alpha_i|\alpha_j)} \Omega_j(l+1) \Omega_i(k)$$

$$x_{i,k+1}^- x_{jl}^- - q^{-(\alpha_i|\alpha_j)} x_{jl}^- x_{i,k+1}^- = q^{-(\alpha_i|\alpha_j)} x_{ik}^- x_{j,l+1}^- - x_{j,l+1}^- x_{ik}^-$$

Proposition

For a quantum affine algebra associated to any untwisted affine Lie algebra, there exists a unique non-degenerate symmetric form $(-, -)$ defined on \mathcal{N}_q^- satisfying $(x_{ij}^- a, b) = (a, \Omega_i(-j)b)$ and $(1, 1) = 1$. Moreover, \mathcal{N}_q^- is a simple left \mathcal{K}_q -module such that

$$\mathcal{N}_q^- \cong \mathcal{K}_q / \left(\sum_{i=1}^N \sum_{k \in \mathbb{Z}} \mathcal{K}_q \Omega_i(k) \right)$$

Remark: 1) For Lie algebras of type ADE the result was shown by Cox, F., Misra, 2015.

2) This allows to define the Kashiwara-like operators on imaginary Verma modules.

- We say that a monomial $x_{i_1 k_1}^- \cdots x_{i_r k_r}^-$ is ordered if $i_1 + k_1 \geq i_2 + k_2 \geq \cdots \geq i_r + k_r$

There exists a product, denoted \star , such that $x_{im}^- \star x_{jn}^-$ is ordered (for ordered monomials: $x_{i_1 k_1}^- \cdots x_{i_l k_l}^- = x_{i_1 k_1}^- \star \cdots \star x_{i_l k_l}^-$)

We define the operator \tilde{x}_{jm}^- on \star -monomials as a \star -left multiplication. We also define the operator $\tilde{\Omega}_i(m)$ on \star -monomials inductively:

$$\tilde{\Omega}_i(m)(x_{jk}^-) := \delta_{ij} \delta_{-m,k},$$

$$\begin{aligned} \tilde{\Omega}_i(m)((x_{i_1 k_1}^- \star (\cdots \star (x_{i_{l-1} k_{l-1}}^- \star x_{i_l k_l}^-) \cdots))) = \\ \delta_{ii_1} \delta_{-m, k_1} (x_{i_2 k_2}^- \star (\cdots \star (x_{i_{l-1} k_{l-1}}^- \star x_{i_l k_l}^-) \cdots)) \end{aligned}$$

$$+ \sum_{r \geq 0} q^{p_{i_1}^{mk_1}} g_{i, i_1, q^{-1}}(r) (x_{i_1, m_1+r}^- \star \tilde{\Omega}_i(m-r)(x_{i_2 k_2}^- \star (\cdots \star (x_{i_{l-1} k_{l-1}}^- \star x_{i_l k_l}^-))))$$

- Define the action of \tilde{X}_{jm}^- and $\tilde{\Omega}_i(m)$ on $\tilde{M}_q(\lambda)$:

$$\tilde{X}_{jm}^-(x_{i_1 k_1}^- \cdots x_{i_r k_r}^- v_\lambda) = \tilde{X}_{jm}^-(x_{i_1 k_1}^- \star \cdots \star x_{i_r k_r}^- v_\lambda) := (x_{jm}^- \star x_{i_1 k_1}^-) \star \cdots \star x_{i_r k_r}^- v_\lambda$$

$$\begin{aligned} & \tilde{\Omega}_i(m)(x_{i_1 k_1}^- \star \cdots \star x_{i_l k_l}^- v_\lambda) \\ &= \delta_{ii_1} \delta_{-m, k_1} x_{i_2 k_2}^- \star \cdots \star x_{i_l k_l}^- v_\lambda \\ &+ \sum_{r \geq 0} q^{p_{i_1}^{mk_1}} g_{i, i_1, q^{-1}}(r) x_{i_1, m_1+r}^- \star \tilde{\Omega}_i(m-r)(x_{i_2 k_2}^- \star \cdots \star x_{i_l k_l}^-) v_\lambda \end{aligned}$$

- Set $\langle x_{im}^-, x_{jn}^- \rangle := (1, \tilde{\Omega}_i(-m)(x_{jn}^-)) \in \mathbb{Z}[q]$

Crystal-like basis

Let $\mathbb{A}_0 = \mathbb{C}[q^{1/2}]_{(q)}$ the ring of rational functions in $q^{1/2}$ regular at 0. Let $\Sigma = \{\mu - k\alpha + n\delta \mid \mu \in P, \alpha \in \Delta^+(\mathfrak{g}), k > 0, n \in \mathbb{Z}\} \cup \{\mu\}$.

• Let M be a $U_q(\hat{\mathfrak{g}})$ -module. We call a free \mathbb{A}_0 -submodule \mathcal{L} of M an *imaginary crystal lattice* of M if the following holds:

1. $\mathbb{C}(q^{1/2}) \otimes_{\mathbb{A}_0} \mathcal{L} \cong M$.
2. $\mathcal{L} \cong \bigoplus_{\lambda \in \Sigma} \mathcal{L}_\lambda$ and $\mathcal{L}_\lambda = \mathcal{L} \cap M_\lambda$.
3. $\tilde{\Omega}_{\psi_i}(m)\mathcal{L} \subseteq \mathcal{L}$ and $\tilde{x}_{im}^-\mathcal{L} \subseteq \mathcal{L}$, for $i \in I$ and $m \in \mathbb{Z}$.

Crystal-like basis

Let $\mathbb{A}_0 = \mathbb{C}[q^{1/2}]_{(q)}$ the ring of rational functions in $q^{1/2}$ regular at 0. Let $\Sigma = \{\mu - k\alpha + n\delta \mid \mu \in P, \alpha \in \Delta^+(\mathfrak{g}), k > 0, n \in \mathbb{Z}\} \cup \{\mu\}$.

• Let M be a $U_q(\hat{\mathfrak{g}})$ -module. We call a free \mathbb{A}_0 -submodule \mathcal{L} of M an *imaginary crystal lattice* of M if the following holds:

1. $\mathbb{C}(q^{1/2}) \otimes_{\mathbb{A}_0} \mathcal{L} \cong M$.
2. $\mathcal{L} \cong \bigoplus_{\lambda \in \Sigma} \mathcal{L}_\lambda$ and $\mathcal{L}_\lambda = \mathcal{L} \cap M_\lambda$.
3. $\tilde{\Omega}_{\psi_i}(m)\mathcal{L} \subseteq \mathcal{L}$ and $\tilde{x}_{im}^- \mathcal{L} \subseteq \mathcal{L}$, for $i \in I$ and $m \in \mathbb{Z}$.

• The \mathbb{A}_0 -module

$$\mathcal{L}(\lambda) := \bigoplus_{\substack{k \geq 0 \\ i_1 + m_1 \geq \dots \geq i_k + m_k \\ i_j, m_j \in \mathbb{Z}}} \mathbb{A}_0 x_{i_1 m_1}^- \cdots x_{i_k m_k}^- v_\lambda$$

is an imaginary crystal lattice of the irreducible reduced imaginary Verma module $\tilde{M}_q(\lambda)$. Moreover,

$$\mathcal{L}(\lambda) = \{u \in \tilde{M}_q(\lambda) \mid \langle u, \mathcal{L}(\lambda) \rangle \subset \mathbb{A}_0\}$$

- An *imaginary crystal base* of an irreducible reduced imaginary Verma module $\tilde{M}_q(\lambda)$ is a pair $(\mathcal{L}, \mathcal{B})$ satisfying:
 1. \mathcal{L} is an imaginary crystal lattice of M .
 2. \mathcal{B} is a \mathbb{C} -basis of $\mathcal{L}/q\mathcal{L} \cong \mathbb{C} \otimes_{\mathbb{A}_0} \mathcal{L}$.
 3. $\mathcal{B} = \cup_{\mu \in \Sigma} \mathcal{B}_\mu$ where $\mathcal{B}_\mu = \mathcal{B} \cap (\mathcal{L}_\mu/q\mathcal{L}_\mu)$.
 4. $\tilde{x}_{im}^- \mathcal{B} \subset \pm \mathcal{B} \cup \{0\}$ and $\tilde{\Omega}_i(m) \mathcal{B} \subset \pm \mathcal{B} \cup \{0\}$.
 5. For $m \in \mathbb{Z}$ and $i \in I_0$ if $\tilde{\Omega}_i(-m)b \neq 0$ and $\tilde{x}_{im}^- b \neq 0$ for $b \in \mathcal{B}$, then $\tilde{x}_{im}^- \tilde{\Omega}_i(-m)b = \tilde{\Omega}_i(-m)\tilde{x}_{im}^- b$.

- An *imaginary crystal base* of an irreducible reduced imaginary Verma module $\tilde{M}_q(\lambda)$ is a pair $(\mathcal{L}, \mathcal{B})$ satisfying:
 1. \mathcal{L} is an imaginary crystal lattice of M .
 2. \mathcal{B} is a \mathbb{C} -basis of $\mathcal{L}/q\mathcal{L} \cong \mathbb{C} \otimes_{\mathbb{A}_0} \mathcal{L}$.
 3. $\mathcal{B} = \cup_{\mu \in \Sigma} \mathcal{B}_\mu$ where $\mathcal{B}_\mu = \mathcal{B} \cap (\mathcal{L}_\mu/q\mathcal{L}_\mu)$.
 4. $\tilde{x}_{im}^- \mathcal{B} \subset \pm \mathcal{B} \cup \{0\}$ and $\tilde{\Omega}_i(m) \mathcal{B} \subset \pm \mathcal{B} \cup \{0\}$.
 5. For $m \in \mathbb{Z}$ and $i \in I_0$ if $\tilde{\Omega}_i(-m)b \neq 0$ and $\tilde{x}_{im}^- b \neq 0$ for $b \in \mathcal{B}$, then $\tilde{x}_{im}^- \tilde{\Omega}_i(-m)b = \tilde{\Omega}_i(-m)\tilde{x}_{im}^- b$.
- For $\lambda \in \hat{\mathfrak{h}}^*$ define

$$\mathcal{B}(\lambda) = \left\{ x_{i_1 m_1}^- \cdots x_{i_k m_k}^- v_\lambda + q\mathcal{L}(\lambda) \in \mathcal{L}(\lambda)/q\mathcal{L}(\lambda) \right. \\ \left. \left| \begin{array}{l} i_1 + m_1 \geq \cdots \geq i_k + m_k, \\ m_1, \dots, m_k \in \mathbb{Z}, i_1, \dots, i_k \in I_0 \end{array} \right. \right\}$$

Theorem [Arias, V.F., Misra, 2024]

If $\lambda \in \hat{\mathfrak{h}}^*$ such that $\lambda(c) = 0$ and $\lambda(h_i) \neq 0$ for all $i \in I_0$, then $(\mathcal{L}(\lambda), \mathcal{B}(\lambda))$ is an imaginary crystal base for $\tilde{M}_q(\lambda)$.

Theorem [Arias, V.F., Misra, 2024]

If $\lambda \in \hat{\mathfrak{h}}^*$ such that $\lambda(c) = 0$ and $\lambda(h_i) \neq 0$ for all $i \in I_0$, then $(\mathcal{L}(\lambda), \mathcal{B}(\lambda))$ is an imaginary crystal base for $\tilde{M}_q(\lambda)$.

• Let $\hat{\mathfrak{h}}_{red}^* = \{\lambda \in P \mid \lambda(c) = 0, \lambda(h_i) \neq 0, i \in I_0\}$. Consider the category $\mathcal{O}_{red,im}^q$ of $U_q(\hat{\mathfrak{g}})$ -modules M such that

1. M is a weight module with weights in $\hat{\mathfrak{h}}_{red}^*$.
2. For any $i \in I_0$ and any $n \in \mathbb{Z}$, x_{in}^+ is locally nilpotent.
3. M is G_q -compatible.

We have in $\mathcal{O}_{red,im}^q$ (Arias, V.F., Oliveira, 2025):

- (1) If $\lambda, \mu \in \hat{\mathfrak{h}}_{red}^*$ then $\text{Ext}_{\mathcal{O}_{red,im}^q}^1(\tilde{M}_q(\lambda), \tilde{M}_q(\mu)) = 0$.
- (2) If M is an irreducible module in the category $\mathcal{O}_{red,im}^q$, then $M \cong \tilde{M}_q(\lambda)$ for some $\lambda \in \hat{\mathfrak{h}}_{red}^*$. Moreover, if N is an arbitrary object of $\mathcal{O}_{red,im}^q$ then $N \cong \bigoplus_{\lambda_i \in \hat{\mathfrak{h}}_{red}^*} \tilde{M}(\lambda_i)$, for some λ_i 's.

Theorem [Arias, V.F., Misra, 2024]

The operators \tilde{x}_{in}^- and $\tilde{\Omega}_i(m)$ are well defined on objects of the category $\mathcal{O}_{red,im}^q$. Every object of the category has an imaginary crystal basis.

Remark: For $\mathfrak{g} = sl(2)$ the result was shown by Cox, F., Misra, 2017.

Thank you!